arXiv:1009.0695v2 [math-ph] 17 Aug 2011

Rate of convergence of linear functions on the unitary group

J. P. Keating, F. Mezzadri and B. Singphu

Abstract

We study the rate of convergence to a normal random variable of the real and imaginary parts of $\operatorname{Tr} A_N U$, where U is an $N \times N$ random unitary matrix and A_N is a deterministic complex matrix. We show that the rate of convergence is $O\left(N^{-2+b}\right)$, with $0 \le b < 1$, depending only on the asymptotic behaviour of the singular values of A_N; for example, if the singular values are non-degenerate, different from zero and $O(1)$ as $N \to \infty$, then $b = 0$. The proof uses a Berry-Esséen inequality for linear combinations of eigenvalues of random unitary matrices, and so appropriate for strongly dependent random variables.

2010 MSC: 15B52, 60F05

1 Introduction

The value distributions of traces of random unitary matrices have been studied extensively over the past fifteen years [9, 13, 15, 24, 11, 14, 21, 25]. The main reason is that they are connected with the linear statistics

$$S_N(\chi) := \chi\left(e^{i\theta_1}\right) + \cdots + \chi\left(e^{i\theta_N}\right), \tag{1.1}$$

where χ is a suitable test function and $e^{i\theta_1}, \ldots, e^{i\theta_N}$ are the eigenvalues of $N \times N$ unitary matrices U distributed according to Haar measure. It turns out that in many applications in particle physics, open quantum systems, quantum chromodynamics and scattering theory it is interesting to understand the asymptotic ($N \to \infty$) behaviour not only of $\operatorname{Tr} U$ but of the more general random variable

$$Z_N := V_N + iW_N = \operatorname{Tr} A_N U, \tag{1.2}$$

where V_N (respectively W_N) is the real (imaginary) part of Z_N, and A_N is a deterministic complex matrix. (See, *e.g.*, [22, 2, 3, 23] and references therein.) In other words, we want to understand the distribution of linear combinations of the elements of random unitary matrices. In general, this type of question arises when Random Matrix Theory is applied to non-Hermitian quantum mechanics, an area of physics which has grown rapidly in the last decades (see, *e.g.*, [19, 20] and references therein). As we shall see, the invariance of Haar measure on $\mathrm{U}(N)$ under group action implies that the distributions of V_N and W_N are the same. Therefore, we shall restrict our attention to V_N.

Samuel [22] and Bars [2] computed the first few terms in the cumulant expansion of V_N, which implicitly show that it converges in distribution to a normal random variable when $N \to \infty$. D'Aristotile *et al.* [10] gave a rigorous proof of this result. Collins and Stolz [6] proved a multivariate version of this theorem: they showed that a vector of the form

$$\left(\operatorname{Tr} A_N^{(1)} U, \ldots, \operatorname{Tr} A^{(r)} U\right), \tag{1.3}$$

where r is independent of N, converges to a joint normal distribution.

In her PhD thesis, Meckes [17, 18] studied the rate of convergence of V_N to a central limit theorem using Stein's method of exchangeable pairs. Let us normalise A_N so that $\operatorname{Tr} A_N A_N^* = N$, where A_N^* is the conjugate transpose of A_N, and denote by $\mathcal{N}(\mu, \sigma^2)$ a normal random variable with mean μ and variance σ^2. Meckes proved that the distance of V_N to $\mathcal{N}(0, 1/2)$ in the total variation metric on probability measures is bounded by $c_N N^{-1}$, where c_N is asymptotic to $2\sqrt{2}$. Chatterjee and Meckes [5] obtained a rate of order $O(N^{-1})$ in the multivariate setting too, and showed that the constant is linear in r.

The bound computed by Meckes holds for any $A_N \in \mathbb{C}^{N \times N}$, subject to the constraint $\operatorname{Tr} A_N A_N^* = N$. However, given a fixed sequence $\{A_N\}_{N>1}$, it is natural to ask how the rate of convergence of V_N depends on A_N. The purpose of this paper is to show that this rate is $O\left(N^{-2+b}\right)$, where $0 \le b < 1$, depending only on the leading order asymptotics as $N \to \infty$ of the greatest singular value of A_N. For example, if the elements of A_N do not grow with N — which is what one would expect for a generic sequence $\{A_N\}_{N>1}$ — then $b = 0$ and the rate of convergence is $O\left(N^{-2}\right)$. When $b = 1$ only a finite fraction of the singular values is different from zero in the limit $N \to \infty$. For technical reasons, which we will discuss in section 3.2, we exclude the case $b = 1$. Meckes' bound $c_N N^{-1}$ does not discern the dependence of the rate of convergence on the singular values of A_N, and our result implies that it is sharp only when $b = 1$.

Our approach is based on the method of moments, which allows us to prove a Berry-Esséen inequality for the eigenvalues of random unitary matrices. In general, Berry-Esséen bounds are used to prove central limit theorems for sums of independent or weakly dependent random variables. It is notable that such a bound exists for sums of eigenvalues of matrices in $\mathrm{U}(N)$, which are strongly correlated.

When A_N is the identity, then Z_N is a class function and the underlying group structure of $\mathrm{U}(N)$ can be exploited. For general A_N these group-theoretical tools are not available. There is a considerable literature addressing the problem of the distribution of $\operatorname{Tr} U^j$, where $j \in \mathbb{Z}_+$. Diaconis and Shahshahani [9], and independently Haake *et al.* [13], proved that it convergences in distribution to $\sqrt{j} Z$, where Z is a standard normal complex random variable. Diaconis and Shahshahani's proof is based on the method of moments; they showed that the k-th moments of $\operatorname{Tr} U^j / \sqrt{j}$ are exactly Gaussian for $k \le N$. This property prompted Diaconis to conjecture that the convergence to a normal random variable is very fast, either exponential or even superexponential. Consider the error

$$e(N) := \sup_{x\in\mathbb{R}} \left|F_N(x) - \Phi(x)\right|, \tag{1.4}$$

where

$$\Phi(x) := \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{x} e^{-\frac{t^2}{2}} dt \tag{1.5}$$

and $F_N(x)$ is the distribution function of $\sqrt{2/j}\,\operatorname{Re}\operatorname{Tr} U^j$, *i.e.*

$$F_N(x) := \int_{-\infty}^{x} f_N\left(t\right) dt, \tag{1.6}$$

where f_N is the probability density function (*p.d.f.*). Johannson [15] proved that $e(N) = O_\epsilon\left(N^{-\epsilon N}\right)$. He also showed that the distance of $\sqrt{2/j}\,\operatorname{Re}\operatorname{Tr} U^j$ to $\mathcal{N}(0,1)$ in the total variation norm is of the same order. Such a rate of convergence to a central limit theorem is unusual in probability theory. The approach that we use to achieve our bounds also sheds light on why the convergence of $\operatorname{Tr} U^j$ is so fast.

Subsequently, many authors have refined or improved Diaconis and Shahshahani's results. Soshnikov [24] showed that the linear statistics (1.1) converge in distribution to a normal random variable in the *mesoscopic* regime too, *i.e.* if one considers eigenvalues in an arc of length L_N with $L_N/N \to 0$ as $N \to \infty$. Hughes and Rudnick [14] studied the *scaling limit* $L_N = N$. It turns out that the number of moments of $S_N(\chi)$ that are exactly Gaussian depends on the class of test functions considered. Diaconis and Evans [11] used the results in [9] to study the asymptotic distributions of integrals of the type $\int_{\mathbb{S}} f d\Xi$, where $\mathbb{S}$ is the unit circle and Ξ_N is the random point measure that places a unit mass at each eigenvalue $e^{i\theta_j}$. Pastur and Vasilchuk [21] and Stolz [25] gave alternative proofs of the convergence to normal random variables of $\operatorname{Tr} U^j$.

This article is structured as follows. In §2 we discuss the background of the problem and introduce our main results. The moments and cumulants of V_N can be computed using the character theory of the symmetric group; these calculations are detailed in §3. In §4 we present the proof of the Berry-Esséen inequality. Finally, §5 and §6 are devoted to the proofs of the main theorems.

2 Statement of results

2.1 Preliminaries

Let us introduce the random variables

$$X_N := \left(\operatorname{Re}\operatorname{Tr} A_N U\right)/\sigma = V_N/\sigma \quad \text{and} \quad Y_N := \left(\operatorname{Im}\operatorname{Tr} A_N U\right)/\sigma = W_N/\sigma, \tag{2.1}$$

where U is an $N \times N$ unitary matrix distributed according to Haar measure and

$$\sigma^2 := \frac{\operatorname{Tr} A_N A_N^*}{2N}. \tag{2.2}$$

The matrices in a given sequence $\{A_N\}_{N>1}$ can be normalised so that σ^2 is independent of N.

Using the polar decomposition we can factorize A_N in the product

$$A_N = H_N V, \tag{2.3}$$

where $V \in \mathrm{U}(N)$ and $H_N = \sqrt{A_N A_N^*}$ is positive-semidefinite. Let us also write $U = W\Theta W^*$, where $W \in \mathrm{U}(N)$ and $\Theta = \operatorname{diag}\left(e^{i\theta_1}, \dots, e^{i\theta_N}\right)$. Since Haar measure is invariant under group action, the random variable $\operatorname{Tr} H_N U/\sigma$ has the same distribution as $\operatorname{Tr} A_N U/\sigma$. Thus, without loss of generality, we can restrict A_N to the set of positive-semidefinite matrices. Furthermore, we have

$$\operatorname{Tr} A_N U = \operatorname{Tr} A_N W\Theta W^* = \operatorname{Tr} W^* A_N W \Theta = \operatorname{Tr} \tilde{A}_N \Theta = \sigma \sum_{j=1}^{N} a_j e^{i\theta_j}, \tag{2.4}$$

where $\tilde{A}_N$ is Hermitian positive-semidefinite too and $\sigma a_j \geq 0$ are its diagonal elements. Therefore, we can write

$$X_N = a_1 \cos\theta_1 + \cdots + a_N \cos\theta_N, \tag{2.5a}$$
$$Y_N = a_1 \sin\theta_1 + \cdots + a_N \sin\theta_N. \tag{2.5b}$$

Since Haar measure is invariant under translation, X_N and Y_N have the same probability distribution. Thus, we shall restrict our attention to X_N.

The characteristic function of X_N is defined by

$$\psi_N(\xi) := \mathbb{E}_{\mathrm{U}(N)}\left\{e^{i\xi X_N}\right\}. \tag{2.6}$$

It admits a representation as an integral over the unitary group. We have

$$\psi_N(\xi) = \int_{\mathrm{U}(N)} \exp\left(\frac{i\xi}{2\sigma}\Big(\operatorname{Tr} A_N U + \operatorname{Tr} A_N^* U^*\Big)\right) d\mu_{\mathrm{H}}(U), \tag{2.7}$$

where $d\mu_{\mathrm{H}}$ denotes Haar measure over $\mathrm{U}(N)$. When A_N is not singular, such an integral can be evaluated explicitly [4] (see also [23] when the matrix in the second trace is different from A_N^*):

$$\psi_N(\xi) = \left(\frac{2\sigma}{\xi}\right)^{\frac{N(N-1)}{2}} \left(\prod_{j=1}^{N-1} j!\right) \frac{\det_{N\times N}\left(\nu_j^{k-1} J_{k-1}\left(\xi\nu_j/\sigma\right)\right)}{\prod_{1\leq j<k\leq N}\left(\nu_k^2 - \nu_j^2\right)}, \tag{2.8}$$

where $\nu_1, \dots, \nu_N$ are the singular values of A_N and J_k is the Bessel function of the first kind. Unfortunately, this beautiful formula is not the best starting point for a straightforward asymptotic analysis. In order to determine the rate of convergence of X_N, we will need to control $\psi_N(\xi)$ when ξ grows like a power of N. This means that N appears as a parameter in both the argument and the index of the Bessel functions. The facts that the asymptotic limit of $J_k(x)$ as $x \to \infty$ is not uniform in the index, and that all the Bessel functions from J_0 to J_{N-1} appear in the determinant, render the analysis of formula (2.8) difficult.

Damgaard and Splittorff [7] computed the first few terms of the asymptotic expansions of integral (2.7) for "low-mass" and "large-mass". In our formalism, this means in the limit as $\xi \to 0$ and $\xi \to \infty$.

The approach that we adopt is based on the method of moments, which can be computed explicitly up to the $2N$-th for any matrix A_N, whether singular or not. The only constraint that we impose on the sequence $\{A_N\}_{N>1}$ is the normalisation (2.2).

Our results will depend on the asymptotic properties of the singular values of A_N. Therefore, we need to introduce quantities that characterise their behaviour in the limit as $N \to \infty$. Let us order the singular values of A_N so that $\nu_1 \leq \cdots \leq \nu_N$ and let $\nu_N^2 = O\left(N^b\right)$. Then, define

$$k := \inf\left\{c \in \mathbb{R} \middle|\, \nu_N^2 \leq cN^b,\ \forall N > 1\right\}. \tag{2.9}$$

Since b is optimal, the normalization (2.2) implies that $0 \leq b \leq 1$ and $k > 0$. The meaning of b and k can be illustrated with a few examples. If all the matrix elements of A_N are $O(1)$ as $N \to \infty$, then $b = 0$. Alternatively, consider the sequence of matrices

$$A_N = \operatorname{diag}\left(\sqrt{2N}, 0, \ldots, 0\right). \tag{2.10}$$

Then $b = 1$ and $k = 2$. In other words, b not only gives the rate of growth of ν_N, but also measures how sparse the set of singular values is in the limit $N \to \infty$.

2.2 Rates of convergence

Using the same notation as in §1, F_N and Φ will denote the distribution functions of X_N and of a standard normal random variable respectively; similarly, f_N is the *p.d.f.* Furthermore, we shall write

$$\phi(x) := \frac{e^{-x^2/2}}{\sqrt{2\pi}} \quad \text{and} \quad \psi(\xi) := \int_{-\infty}^{\infty} e^{i\xi x}\phi(x)dx = e^{-\xi^2/2}. \tag{2.11}$$

Theorem 2.1. *Suppose* $\{A_N\}_{N>1}$ *is a sequence of matrices such that* $\sigma^2 = \operatorname{Tr} A_N A_N^*/(2N)$ *is independent of* N *and that* $0 \leq b < 1$*. We have*

$$e(N) := \sup_{x\in\mathbb{R}} |F_N(x) - \Phi(x)| = O\left(N^{-2+b}\right), \quad N \to \infty. \tag{2.12}$$

As we shall see, the power of minus two in (2.12) is determined by the Haar measure on U(N). The sequence $\{A_N\}_{N>1}$ influences the rate of convergence only through the parameter b, which is a measure of the asymptotic distribution of the singular values of the matrices A_N.

We can prove an analogous statement in the total variation norm.

Theorem 2.2. *Let $\{A_N\}_{N>1}$ be a sequence satisfying the same conditions as in theorem* 2.1*. We have*

$$\int_{-\infty}^{\infty} |f_N(x) - \phi(x)|\, dx = O_\epsilon\left(N^{-2+b+\epsilon}\right), \quad N \to \infty, \tag{2.13}$$

where $\epsilon \in \left(0, \frac{1}{2}(1-b)\right)$.

As we discussed in the introduction, for technical reasons theorems 2.1 and 2.2 exclude $b = 1$. Meckes' [18] result suggests that they are correct for $b = 1$ too.

The starting formula to prove theorems 2.1 and 2.2 is

$$e\,(N) \le \frac{1}{\pi}\int_{-T_N}^{T_N} \left|\frac{\psi_N(\xi) - \psi(\xi)}{\xi}\right| d\xi + \frac{24m}{\pi T_N} \tag{2.14}$$

(see [12], p. 538), where

$$m := \max_{x\in\mathbb{R}} |\phi(x)| = \frac{1}{\sqrt{2\pi}} \tag{2.15}$$

and T_N is an appropriate cut-off. Formula (2.14) transfers the problem of computing $e(N)$ into that of finding a bound for $|\psi_N(\xi) - \psi(\xi)|$ for sufficiently large ξ.

Theorem 2.3. *Let C and δ be two constants independent of N and let $0 \le |\xi| < \delta N^{(1-b)/2}$. We have*

$$|\psi_N(\xi) - \psi(\xi)| \le \frac{C\xi^4}{N^{2-b}} e^{-\xi^2/2}. \tag{2.16}$$

Throughout this paper C will denote a constant, which may be different at each occurrence.

Remark 2.4. *Theorem* 2.3 *is of interest in its own right. Such bounds are called **Berry-Esséen inequalities**. They determine rates of convergence to central limit theorems, usually for sums of independent or weakly dependent random variables. The eigenvalues of random unitary matrices, however, exhibit a high degree of correlation.*

For eigenvalues of random unitary matrices, one consequence of such a strong dependence is that the variance $\sigma^2 = \mathrm{Tr}\, A_N A_N^*/(2N)$ remains finite in the limit $N \to \infty$. Instead, the variance of the sum of N independent random variables grows linearly in N. When the moments diverge in the limit $N \to \infty$, just the first few are enough to determine an optimal bound. Since the right-hand sides of equations (2.5a) *and* (2.5b) *converge to normal random variables without any normalisation, the proof of theorem* 2.3 *requires knowing the first $2N$ moments of X_N.*

3 Moments and cumulants of X_N

The purpose of this section is to provide bounds and asymptotic formulae for the moments and cumulants of X_N that will be needed to prove theorem 2.3. Most of these can be derived from the results of Samuel [22], which we summarise in §3.1.

3.1 Averages of matrix elements and the symmetric group

Samuel [22] studied averages of the form

$$\int_{\mathrm{U}(N)} U_{i_1 j_1} \cdots U_{i_m j_m} \overline{U}_{k_1 l_1} \cdots \overline{U}_{k_m l_m} d\mu_{\mathrm{H}}(U) \\ = \sum_{\sigma,\tau \in \mathfrak{S}_m} M_{\sigma,\tau}(N) \delta_{i_1 k_{\sigma 1}} \cdots \delta_{i_m k_{\sigma m}} \delta_{j_1 l_{\tau 1}} \cdots \delta_{j_m l_{\tau m}}, \quad (3.1)$$

where $\mathfrak{S}_m$ denotes the symmetric group of degree m. The moments of X_N are simply linear combinations of these integrals.

All the information on the averages (3.1) is contained in the coefficients $M_{\sigma,\tau}(N)$. A permutation of m letters can always be factorised in a product of cycles. It turns out that $M_{\sigma,\tau}(N)$ depends only on the cycle decomposition of $\sigma\tau^{-1}$.

The lengths of the cycles of a permutation identify a sequence of non-negative integers $\lambda = (\lambda_1, \dots, \lambda_k)$ such that

$$\lambda_1 \geq \cdots \geq \lambda_k \quad \text{and} \quad |\lambda| := \lambda_1 + \cdots + \lambda_k = m. \quad (3.2)$$

In other words, there exists a one-to-one correspondence between the cycle structures of $\mathfrak{S}_m$ and the set of *partitions* of m. The partition $\lambda(g)$ is called *cycle-type* of $g \in \mathfrak{S}_m$. Therefore, we shall adopt the notation

$$M_\lambda(N) := M_{\sigma,\tau}(N), \quad (3.3)$$

where λ is the cycle-type of $\sigma\tau^{-1}$.

A partition of m is denoted by $\lambda \vdash m$; the addends λ_j are the *parts* of λ. An alternative notation for a partition is the *frequency representation*: if λ contains r_1 1s, r_2 2s and so forth, we write $\lambda = (1^{r_1} 2^{r_2} \dots m^{r_m})$. The length of a partition $\ell(\lambda)$ is the largest j such that $\lambda_j > 0$. We also have

$$\ell(\lambda) = r_1 + \cdots + r_m. \quad (3.4)$$

We shall find it convenient not to distinguish between two partitions that differ only by a sequence of zeros at the end. For example, $(3,1,1)$ and $(3,1,1,0,0,0)$ are clearly the same partition.

Elements of $\mathfrak{S}_m$ that belong to the same conjugacy class share the same cycle-type. Therefore, the conjugacy classes of $\mathfrak{S}_m$ can be labelled by the set of the partitions of m. The number of elements in the conjugacy class λ is

$$g_\lambda := \frac{m!}{1^{r_1} r_1! \cdots m^{r_m} r_m!}. \quad (3.5)$$

Furthermore, the conjugacy classes of $\mathfrak{S}_m$ are in one-to-one correspondence with its irreducible representations, which can be identified with the set of partitions of m too. Since characters are class functions they depend only on the cycle-types of the permutations.

The notation χ^{μ}_{λ} indicates the character of the irreducible representation μ evaluated on elements of cycle-type λ.

Sometimes it is convenient to represent partitions using Young tableaux. If $\lambda = (\lambda_1, \ldots, \lambda_k)$, we draw k left-justified rows of boxes, or nodes; the top row should contain λ_1 boxes, the next one λ_2 and so on. For example, let $\lambda = (5, 4, 4, 3, 1)$. Then,



is the corresponding Young tableau.

Samuel [22] derived an explicit formula for $M_\lambda(N)$ when $m \le N$:

$$M_\lambda(N) := \frac{1}{m!} \sum_{\mu \vdash m} \frac{\dim V_\mu}{f_\mu(N)} \chi^{\mu}_{\lambda}, \quad \lambda \vdash m, \tag{3.6}$$

where

$$f_\lambda(N) := \frac{1}{\dim V_\lambda} \sum_{\mu \vdash m} g_\mu \chi^{\lambda}_{\mu} N^{\ell(\mu)} \tag{3.7}$$

and

$$\dim V_\lambda = m! \frac{\prod_{1 \le i < j \le \ell(\lambda)} (\lambda_i - \lambda_j + j - i)}{\prod_{j=1}^{\ell(\lambda)} (\lambda_j + \ell(\lambda) - j)} \tag{3.8}$$

is the dimension of the irreducible representation V_λ.

The right-hand side of (3.7) is polynomial in N of degree m. It turns out that $f_\lambda(N)$ has only integer roots, which have a simple representation in terms of the Young tableau of λ; they are given by all the differences $i - j$, where i counts the rows of the diagram in descending order and j counts its columns from left to right. For example, if $\lambda = (5, 4, 4, 3, 1)$, then the roots of $f_\lambda(N)$ are

0	- 1	-2	- 3	-4
1	0	-1	-2	
2	1	0	-1	
3	2	1		
4				

We shall give a proof of this property later in this section.

Since the characters of the irreducible representations of the symmetric group are know via Frobenius's character formula, equations (3.5), (3.6) and (3.7) completely determine the averages (3.1).

Let $\lambda_1, \ldots, \lambda_k$ be the parts of a partition $\lambda \vdash m$. (We do not impose any ordering on the λ_js.) The coefficients $M_\lambda(N)$ obey the recursion relations [22]

$$\begin{aligned}\delta_{\lambda_k 1} M_{(\lambda_1,\ldots,\lambda_{k-1})}(N) = N M_{(\lambda_1,\ldots,\lambda_k)}(N) + \sum_{p+q=\lambda_k} M_{(\lambda_2,\ldots,\lambda_k,p,q)}(N) \\ + \sum_{j=1}^{k-1} \lambda_j M_{(\lambda_1,\ldots,\lambda_{j-1},\lambda_j+\lambda_k,\lambda_{j+1},\ldots,\lambda_{k-1})}(N)\,,\end{aligned} \tag{3.9}$$

with initial condition $M_0(N) = 1$. These equations do not depend on permutations of the λ_js and are a complete set, which uniquely determines the coefficients $M_\lambda(N)$ for $\lambda \vdash m$ in terms of those for $\lambda \vdash m-1$.

Traces of powers of matrices are homogeneous symmetric polynomials in the eigenvalues. Symmetric functions are intertwined with the character theory of the symmetric group. Therefore, it is not a surprise that the formalism of symmetric polynomials will become useful in computing the moments and cumulants of X_N.

For every j the *power sum* of m variables is

$$p_j(x_1, \ldots, x_m) := x_1^j + \cdots + x_m^j. \tag{3.10}$$

Next, we extend the definition (3.10) by taking the product

$$p_\lambda := p_{\lambda_1} \cdots p_{\lambda_k} = \prod_{j=1}^m p_j^{r_j}, \tag{3.11}$$

where the r_js are the frequencies of λ. Now suppose that $\ell(\lambda) \le m$. The *Schur function* $s_\lambda(x_1, \ldots, x_m)$ is defined by the ratio of two $m \times m$ determinants:

$$s_\lambda(x_1, \ldots, x_m) := \frac{\det\left(x_i^{\lambda_j + m - j}\right)}{\det\left(x_i^{m-j}\right)}. \tag{3.12}$$

Schur functions are homogeneous symmetric polynomials of degree $|\lambda|$ and are related to the power sums by the formulae (see [16], p.114)

$$s_\lambda = \frac{1}{m!} \sum_{\mu \vdash m} g_\mu \chi_\mu^\lambda p_\mu \quad \text{and} \quad p_\mu = \sum_{\lambda \vdash m} \chi_\mu^\lambda s_\lambda. \tag{3.13}$$

If $x_1, \ldots, x_m$ are the eigenvalues of a $m \times m$ matrix X we write $p_\lambda(X) = p_\lambda(x_1, \ldots, x_m)$ and $s_\lambda(X) = s_\lambda(x_1, \ldots, x_m)$.

Corollary 3.1. *The polynomial* (3.7) *can be factorised as*[1]

$$f_\lambda(N) = \frac{1}{\dim V_\lambda} \sum_{\mu \vdash m} g_\mu \chi_\mu^\lambda N^{\ell(\mu)} = \prod_{(i,j)} (N - i + j) = \prod_{j=1}^{\ell(\lambda)} \frac{(N + \lambda_j - j)!}{(N-j)!}, \tag{3.14}$$

where the pair (i,j) *span the row and column indices of the Young tableau of* λ.

---

[1] Samuel conjectured this formula in the appendix of his article [22], but did not give a proof.

Proof. Let $\lambda \vdash m$ and N be a positive integer, then $p_\lambda(I_N) = N^{\ell(\lambda)}$. Therefore, from formula (3.13) for $N \in \mathbb{Z}_+$

$$f_\lambda(N) = m! \frac{s_\lambda(I_N)}{\dim V_\lambda}. \tag{3.15}$$

The irreducible representations of the symmetric group and of $\mathrm{GL}(N, \mathbb{C})$ are related by the Schur-Weyl duality. If $N \geq \ell(\lambda)$ the Schur functions are precisely the irreducible characters of $\mathrm{GL}(N, \mathbb{C})$. Thus, equation (3.15) connects the dimensions of irreducible representations of $\mathfrak{S}_m$ and $\mathrm{GL}(N, \mathbb{C})$ corresponding to the same λ. Now, we have

$$s_\lambda(I_N) = \prod_{1 \leq j < k \leq N} \frac{\lambda_j - \lambda_k + k - j}{k - j}. \tag{3.16}$$

Combining this formula with (3.15) and (3.8) gives equation (3.14). □

3.2 The moments

Formula (2.7) implies that ψ_N is an entire function. Therefore, the series

$$\psi_N(\xi) = \sum_{n=0}^{\infty} \frac{(i\xi)^n}{n!} \mu_n \tag{3.17}$$

converges in all the complex plane and defines all the moments of X_N, which identify its probability distribution uniquely.

Now, consider the Taylor expansion of the integral (2.7):

$$\psi_N(\xi) = \sum_{n=0}^{\infty} \frac{1}{n!} \left(\frac{i\xi}{2\sigma}\right)^n \sum_{m=0}^{n} \binom{n}{m} \int_{\mathrm{U}(N)} (\mathrm{Tr}\, A_N U)^{n-m} (\mathrm{Tr}\, A_N^* U^*)^m \, d\mu_{\mathrm{H}}(U). \tag{3.18}$$

Since Haar measure is left and right invariant, the integral in this sum is zero unless $n = 2m$. Therefore,

$$\psi_N(\xi) = \sum_{m=0}^{\infty} \frac{(-1)^m}{(m!)^2} \left(\frac{\xi}{2\sigma}\right)^{2m} I_N^m(A_N), \tag{3.19}$$

where

$$I_N^m(A_N) := \int_{\mathrm{U}(N)} |\mathrm{Tr}\, A_N U|^{2m} \, d\mu_{\mathrm{H}}(U). \tag{3.20}$$

Thus, the moments of X_N are given by the formula

$$\mu_{2m} = \frac{(2m-1)!!}{m! \, (2\sigma^2)^m} I_N^m(A_N). \tag{3.21}$$

Proposition 3.2 (Samuel 1980)**.** *Let $m \leq N$ and let $\lambda = (1^{r_1} \ldots m^{r_m})$ denote a partition of m. We have*

$$I_N^m(A_N) = m! \sum_{\lambda \vdash m} g_\lambda M_\lambda(N) \, p_\lambda(A_N A_N^*). \tag{3.22}$$

Proof. The right-hand side of equation (3.1) can be re-written as

$$\sum_{\rho,\tau\in\mathfrak{S}_m} M_\rho\left(N\right)\delta_{i_1 k_{\rho\tau 1}}\cdots\delta_{i_m k_{\rho\tau m}}\delta_{j_1 l_{\tau 1}}\cdots\delta_{j_m l_{\tau m}}, \tag{3.23}$$

where we have shifted the index in the sum by setting $\rho=\sigma\tau^{-1}$ and used the fact that $M_{\sigma,\tau}(N)$ depends only on $\sigma\tau^{-1}$. By multiplying equation (3.23) by $A_{Nj_1i_1},\dots,A_{Nj_mi_m}$ and $\overline{A}_{Nl_1k_1},\dots,\overline{A}_{Nl_mk_m}$ and summing over all indices, we obtain an expression of the form

$$\sum_{\rho\in\mathfrak{S}_m} M_\rho\left(N\right)\sum_{\tau\in\mathfrak{S}_m}\sum_{\dots,\alpha,\beta,\gamma,\delta,\eta,\dots}\cdots A_{N\alpha\beta}\overline{A}_{N\gamma\beta}A_{N\gamma\delta}\overline{A}_{N\eta\delta}\cdots \tag{3.24}$$

Consecutive indices in the inner sum, say β and γ, are of the type $k_{\rho\tau v}$ and $l_{\tau w}$ respectively, where $w=\tau^{-1}\rho\tau v$. Thus, the collection of the addends such that $v=\left(\tau^{-1}\rho\tau\right)^j v$ contributes with a factor $\operatorname{Tr}\left(A_N A_N^*\right)^j$.

Each letter belonging to a cycle of length j is a fixed point of order j of every element in the conjugacy class of ρ. The inner sum in (3.24) depends only on powers of $\tau^{-1}\rho\tau$, and therefore is a class function and is independent of τ. Each cycle of length j produces the factor $\operatorname{Tr}\left(A_N A_N^*\right)^j$. Therefore, we have

$$\sum_{\dots,\alpha,\beta,\gamma,\delta,\eta,\dots}\cdots A_{N\alpha\beta}\overline{A}_{N\gamma\beta}A_{N\gamma\delta}\overline{A}_{N\eta\delta}\cdots=\prod_{j=1}^{m}\left(\operatorname{Tr}\left(A_N A_N^*\right)^j\right)^{r_j}=p_\lambda\left(A_N A_N^*\right). \tag{3.25}$$

Finally, formula (3.22) follows from the fact that $M_\rho\left(N\right)$ is a class function. □

Remark 3.3. *The integral* (3.22)*, and thus by* (3.21) *the moments too, are linear combinations of the coefficients* $M_\lambda(N)$*, which have poles at the zeros of* $f_\lambda(N)$*. Such poles are related to certain singular integrals over* $\mathrm{U}(N)$*, which appear in lattice Quantum Chromodynamics and were first noted by De Wit and 't Hooft* [8]*. They observed that such integrals are divergent for certain values of* N*. The moments of* X_N*, however, are always finite. The reason why the De Wit-'t Hooft anomalies do not affect formula* (3.22) *is because it is correct only for* $m\le N$*, and by corollary* 3.1 *the greatest zero of* $f_\lambda(N)$ *is* $m-1$*.*

As we mentioned at the beginning of this section, in order to prove the Berry-Esséen inequality (2.16) we need bounds and asymptotic formulae for the moments and cumulants. The evaluation of the right-hand side of equation (3.22) requires Frobenius's character formula, which is quite cumbersome to use when explicit formulae are needed. It turns out that (3.22) can be expressed in terms of Schur functions, which allow it to be manipulated explicitly.

Corollary 3.4. *We have*

$$\mu_{2m}=\frac{(2m-1)!!}{\left(2\sigma^2\right)^m}\sum_{\lambda\vdash m}\left(\prod_{j=1}^{\ell(\lambda)}\frac{(N-j)!}{(N+\lambda_j-j)!}\right)\dim V_\lambda\, s_\lambda\left(A_N A_N^*\right). \tag{3.26}$$

Proof. From formulae (3.6), (3.13) and (3.22) we obtain

$$\begin{aligned} I_N^m(A_N) &:= m! \sum_{\lambda \vdash m} g_\lambda M_\lambda(N) p_\lambda\left(A_N A_N^*\right) \\ &= \sum_{\lambda \vdash m} \sum_{\mu \vdash m} \frac{\dim V_\mu}{f_\mu(N)} g_\lambda \chi_\lambda^\mu p_\lambda\left(A_N A_N^*\right) \\ &= m! \sum_{\mu \vdash m} \frac{\dim V_\mu}{f_\mu(N)} s_\mu\left(A_N A_N^*\right). \end{aligned} \tag{3.27}$$

Equation (3.26) follows from formula (3.14). ☐

We are now in a position to find asymptotic formulae for the first N moments of X_N. Let us denote the moments of $\mathcal{N}(0,1)$ by μ_{2m}^{G}, *i.e.*

$$\mu_{2m}^{\mathrm{G}} := (2m-1)!!. \tag{3.28}$$

Proposition 3.5. *We have the following bounds:*

$$\mu_{2m} = \mu_{2m}^{\mathrm{G}}\Big(1 + O\left((mk)^m N^{-1}\right)\Big) \tag{3.29}$$

and

$$\mu_{2m} \le \left(N^b k\right)^m \mu_{2m}^{\mathrm{G}}. \tag{3.30}$$

Proof. From equations (3.6) and (3.22) we have

$$\mu_{2m} = \frac{\mu^{\mathrm{G}}}{\left(2\sigma^2\right)^m m!} \sum_{\lambda \vdash m} \sum_{\mu \vdash m} \frac{\dim V_\mu}{f_\mu(N)} g_\lambda \chi_\lambda^\mu p_\lambda\left(A_N A_N^*\right). \tag{3.31}$$

The first step consists in finding bounds for $p_\lambda\left(A_N A_N^*\right)$ and $f_\mu(N)$. Remember that by definition (2.9), the greatest singular value of A_N is bounded by $\sqrt{kN^b}$, where $k = O(1)$ and $0 \le b < 1$. We have

$$\operatorname{Tr}\left(A_N A_N^*\right)^j = \left(2\sigma^2\right)^j N^{a+bj} c_j\left(A_N\right), \tag{3.32}$$

where $c_j\left(A_N\right) = O\left(k^j\right)$ and $a = 1 - b$. Note that by definition σ is independent of A_N, therefore $c_1\left(A_N\right) = 1$. It follows that

$$p_\lambda\left(A_N A_N^*\right) = \prod_{j=1}^m \left(\operatorname{Tr}\left(A_N A_N^*\right)^j\right)^{r_j} = \left(2\sigma^2\right)^m N^{a\ell(\lambda) + bm} c_\lambda\left(A_N\right), \tag{3.33}$$

where

$$c_\lambda\left(A_N\right) = \prod_{j=1}^m c_j\left(A_N\right)^{r_j} = O\left(k^m\right). \tag{3.34}$$

Denote by $\lambda_e = (1^m)$ the cycle-type of the identity in $\mathfrak{S}_m$. Combining equations (3.33) and (3.34) we obtain.

$$p_\lambda\left(A_N A_N^*\right) = \left(2\sigma^2\right)^m N^m \times \begin{cases} 1 & \text{if } \lambda = \lambda_e, \\ O\big(k^m N^{-a(m-\ell(\lambda))}\big) & \text{if } \lambda \neq \lambda_e. \end{cases} \tag{3.35}$$

Now consider $f_\mu(N)$. We can easily see that for $N \geq m+1$

$$f_{(1^m)}(N) = \prod_{j=1}^{m} (N-j) \leq f_\mu(N) \leq f_{(m^1)}(N) = \prod_{j=1}^{m} (N+j)\,, \quad \mu \vdash m, \tag{3.36}$$

where (m^1) and (1^m) correspond to the trivial and alternating representations respectively, which are both one-dimensional. We can re-write the inequalities (3.36) in the following way

$$\begin{gathered} N^m \prod_{j=1}^{m} \left(1 - \frac{j}{N}\right) \leq f_\mu(N) \leq N^m \prod_{j=1}^{m} \left(1 + \frac{j}{N}\right) \\ \frac{N^m m!}{(m+1)^m} \leq f_\mu\left(N\right) \leq \frac{N^m\left(2m+1\right)\cdots(m+2)}{(m+1)^m}. \end{gathered} \tag{3.37}$$

The sum (3.31) can be split as follows:

$$\begin{aligned} \mu_{2m} = {} & \frac{\mu^{\mathrm{G}} N^m}{m!} \sum_{\mu \vdash m} \frac{\left(\dim V_\lambda\right)^2}{f_\mu(N)} \\ & + \frac{\mu^{\mathrm{G}}}{\left(2\sigma^2\right)^m m!} \sum_{\substack{\lambda \vdash m \\ \lambda \neq \lambda_e}} \sum_{\mu \vdash m} \frac{\dim V_\mu}{f_\mu(N)} g_\lambda \chi^\mu_\lambda p_\lambda\left(A_N A_N^*\right). \end{aligned} \tag{3.38}$$

The first sum on the right-hand side can be estimated using the bounds (3.36)

$$\begin{aligned} \frac{\mu^{\mathrm{G}} N^m}{m!} \sum_{\mu \vdash m} \frac{\left(\dim V_\lambda\right)^2}{f_\mu(N)} &= \frac{\mu^{\mathrm{G}}}{m!} \sum_{\mu \vdash m} \left(\dim V_\lambda\right)^2 \left(1 + O\left(e^m N^{-1}\right)\right) \\ &= \mu^{\mathrm{G}}\big(1 + O\left(e^m N^{-1}\right)\big). \end{aligned} \tag{3.39}$$

Using the same ideas, we write

$$\frac{\mu^{\mathrm{G}}}{\left(2\sigma^2\right)^m m!} \sum_{\mu \vdash m} \chi^\mu_\lambda \frac{\dim V_\lambda}{f_\mu(N)} = \frac{\mu^{\mathrm{G}}}{\left(2\sigma^2\right)^m m! N^m} \sum_{\mu \vdash m} \chi^\mu_\lambda \dim V_\lambda \big(1 + O\left(e^m N^{-1}\right)\big). \tag{3.40}$$

Irreducible representations of finite groups can always be chosen to be unitary. Therefore, we have that $|\chi^\mu_\lambda| \leq \chi^\mu_{\lambda_e} = \dim V_\mu$. Thus, using the orthogonality of the characters, the sum (3.40) becomes

$$\frac{\mu^{\mathrm{G}} R(m,N)}{\left(2\sigma^2\right)^m N^m}, \tag{3.41}$$

where $R(m, N) = O\left(e^m N^{-1}\right)$. Finally, inserting equation (3.41) into (3.38) and using (3.35) we obtain formula (3.29)

In order to prove equation (3.30), recall that from formulae (3.26) and (3.14) we can write

$$\mu_{2m} = \frac{\mu_{2m}^{\mathrm{G}}}{\left(2\sigma^2\right)^m m!} \sum_{\lambda \vdash m} \left(\dim V_\lambda\right)^2 \frac{s_\lambda\left(A_N A_N^*\right)}{s_\lambda\left(I_N\right)}. \tag{3.42}$$

The greatest singular value of A_N is bounded by $2\sigma^2\sqrt{kN^b}$. Since the eigenvalues of $A_N A_N^*$ are non-negative, $s_\lambda\left(A_N A_N^*\right)$ is positive and

$$s_\lambda\left(A_n A_N^*\right) \leq \left(kN^b\right)^m s_\lambda\left(I_N\right). \tag{3.43}$$

Then, equation (3.30) follows from the orthogonality of the characters. □

An immediate corollary is the convergence in distribution of X_N to $\mathcal{N}(0,1)$ (D'Aristotile *et al.* [10]). For fixed m formula (3.29) gives

$$\lim_{N \to \infty} \mu_{2m} = \mu_{2m}^{\mathrm{G}}. \tag{3.44}$$

The bound (3.30) plays an important role in the proof of the Berry-Esséen inequality (2.16). For $N \gg m$ (3.29) is a better bound; however, it becomes much worse when $m \approx N$. This is an important regime. As we shall see, when $b = 1$ the right-hand side of (3.30) is too large to allow (2.16) to be valid for a range of ξ sufficiently large for our purposes. We believe that the correct bound for μ_{2m} is much smaller than both (3.29) and (3.30). The reason is that the sum

$$\sum_{\lambda \vdash m} \sum_{\mu \vdash m} \frac{\dim V_\mu}{f_\mu(N)} g_\lambda \chi_\lambda^\mu p_\lambda\left(A_N A_N^*\right) \tag{3.45}$$

is characterised by a sequence of cancellations.

Remark 3.6. *It is worth noting that, since the integral on the right-hand side of equation* (3.18) *is zero unless $n = 2m$, the proof of proposition* 3.5. *also demonstrates that the random variable $Z_N = X_N + iY_N = \operatorname{Tr} A_N U$ converges in distribution to a complex normal random variable Z, whose centred moments*[2] *are*

$$\mathbb{E}\left\{Z^m \overline{Z}^n\right\} = \delta_{mn} \sigma^{2m} m!. \tag{3.46}$$

---

[2] Here we have adopted the convention that the real and imaginary parts of a standard complex normal random variable have variance $\frac{1}{2}$. Therefore, if we had studied $\operatorname{Tr} A_N U$, instead of its real and imaginary parts separately, we should have set $\sigma^2 = \operatorname{Tr} A_N A_N^*/N$. This explains the discrepancy of a factor 2^m in the notation used in equations (3.47) and (3.46).

Remark 3.7. *When $A_N = I$ the first N moment are exactly gaussian independently of N. This is a particular case of a more general result proved by Diaconis and Shahshahani* [9] *and can be easily recovered in our formalism. We have*

$$\begin{aligned} I_N^m(I) &= m!\left(2\sigma^2\right)^m \sum_{\lambda\vdash m} g_\lambda M_\lambda(N)\, N^{\ell(\lambda)} \\ &= \left(2\sigma^2\right)^m \sum_{\lambda\vdash m}\sum_{\mu\vdash m} \frac{\chi^\mu_{\lambda_e}\chi^\mu_{\lambda_e}}{f_\mu(N)}\frac{g_\lambda\chi^\mu_\lambda}{\chi^\mu_{\lambda_e}} N^{\ell(\lambda)} \\ &= \left(2\sigma^2\right)^m \sum_{\mu\vdash m}\chi^\mu_{\lambda_e}\chi^\mu_{\lambda_e} = m!\left(2\sigma^2\right)^m, \quad m\le N. \end{aligned} \tag{3.47}$$

3.3 The cumulants

The characteristic function $\psi_N(\xi)$ is entire and by definition $\psi_N(0) = 1$. Therefore, the Taylor series

$$\log\psi_N(\xi) = \sum_{n=1}^{\infty}\frac{(i\xi)^n}{n!}\kappa_n \tag{3.48}$$

converges in a neighbourhood of the origin. The coefficients κ_n are by definition the *cumulants* of X_N and determine uniquely its probability distribution. They are related to the moments by the recurrence relation

$$\kappa_n = \mu_n - \sum_{k=1}^{n-1}\binom{n-1}{k-1}\kappa_k\mu_{n-k}. \tag{3.49}$$

The choice of whether to use the moments or the cumulants depends on the information that one is seeking to extract. It turns out that in the proof of the Berry-Esséen inequality (2.16) we shall need the asymptotic behaviour of both. The purpose of this section is to derive a bound for κ_{2m} for $m \le N$.

Let $\lambda \vdash n$ and define

$$\kappa_\lambda := \kappa_1\cdots\kappa_{\lambda_l} = \prod_{j=1}^{n}\kappa_j^{r_j}, \tag{3.50}$$

where the r_js are the frequencies of the partition λ. There exists an elegant formula (see, *e.g.* [16], pp. 30–31) that expresses the moments as polynomials in the cumulants:

$$\mu_n = \sum_{\lambda\vdash n} c_\lambda\kappa_\lambda, \tag{3.51}$$

where

$$c_\lambda := \frac{n!}{(1!)^{r_1}\,r_1!\cdots(m!)^{r_m}\,r_m!} \tag{3.52}$$

is the number of decompositions of a set of n elements into disjoint subsets containing $\lambda_1, \dots, \lambda_n$ elements. Similarly, equation (3.49) can be solved for the cumulants:

$$\kappa_n = \sum_{\lambda \vdash n} (-1)^{\ell(\lambda)-1} \left(\ell\left(\lambda\right) - 1\right)! c_\lambda \mu_\lambda, \tag{3.53}$$

where

$$\mu_\lambda := \mu_1 \cdots \mu_{\lambda_l} = \prod_{j=1}^{n} \mu_j^{r_j}. \tag{3.54}$$

All the odd moments of X_N are zero, therefore all the odd cumulants are zero too. Thus, (3.51) can be rewritten as

$$\mu_{2m} = (2m-1)!! \sum_{\lambda \vdash m} \frac{c_\lambda \kappa_{2\lambda}.}{\left(1!!\right)^{r_j} \cdots \left((2m-1)!!\right)^{r_m}}, \tag{3.55}$$

where we have used the notation $2\lambda = (2\lambda_1, \dots, 2\lambda_l)$.

The $2m$-th moment of X_N is a polynomial of degree m in the traces $\operatorname{Tr}(A_N A_N^*)^j$; the recursion relations (3.49) imply that the $2m$-th cumulant is also a polynomial of degree m in the same variables. Therefore, we can write

$$\kappa_{2m} = \frac{(2m-1)!!}{\left(2\sigma^2\right)^m} \sum_{\lambda \vdash m} g_\lambda K_\lambda(N) p_\lambda\left(A_N A_N^*\right). \tag{3.56}$$

If we know the asymptotic behaviour of $K_\lambda(N)$, then we can determine that of the cumulants. In turn, the coefficients $K_\lambda(N)$ are related to those of $M_\lambda(N)$.

The union $\lambda \cup \mu$ is defined as the partition whose parts are those of λ and μ arranged in descending order. Cumulants have a combinatorial interpretation in term of partitions of sets; let us define

$$M_\lambda(N) := \sum_{\Lambda} a_\Lambda \prod_{\mu \in \Lambda} K'_\mu(N), \tag{3.57}$$

where Λ runs through all possible distinct decompositions of λ as a union of sub-partitions. The meaning of a_Λ is better explained with an example. Consider the partition $\lambda = (\lambda_1, \lambda_2, \lambda_3)$ and write

$$\begin{aligned} M_{(\lambda_1,\lambda_2,\lambda_3)} = K'_{(\lambda_1,\lambda_2,\lambda_3)} &+ K'_{(\lambda_1)} K'_{(\lambda_2,\lambda_3)} + K'_{(\lambda_2)} K'_{(\lambda_1,\lambda_3)} \\ &+ K'_{(\lambda_3)} K'_{(\lambda_1,\lambda_2)} + K'_{(\lambda_1)} K'_{(\lambda_2)} K'_{(\lambda_3)}. \end{aligned} \tag{3.58}$$

If the λ_1, λ_2 and λ_3 are all different, then each summand in (3.58) is distinct, but if some parts of λ are repeated, this is not the case. For example, let $\lambda = (3,1,1)$, then $(\lambda_2) \cup (\lambda_1, \lambda_3)$ and $(\lambda_3) \cup (\lambda_1, \lambda_2)$ are the same decomposition of λ and

$$K'_{(\lambda_2)} K'_{(\lambda_1,\lambda_3)} = K'_{(\lambda_3)} K'_{(\lambda_1,\lambda_2)}. \tag{3.59}$$

The coefficient a_Λ is precisely such a multiplicity. Computing it is an exercise in elementary combinatorics.

Let $\lambda \vdash m$ and define π_μ to be the number of times that a partition μ appears in the decomposition $\lambda = \bigcup_{\mu \in \Lambda} \mu$. Furthermore, let r_j and s_j^μ denote the frequencies of j in λ and μ respectively. We have

$$a_\Lambda = \prod_{j=1}^m r_j! \prod_{\mu \in \Lambda} \frac{1}{s_j^\mu ! \pi_\mu !}. \tag{3.60}$$

Proposition 3.8. *The coefficients* $K_\lambda(N)$ *and* $K'_\lambda(N)$*, defined in equations* (3.56) *and* (3.57) *respectively, coincide.*

Proof. For the sake of simplicity, let us set $x_j := \operatorname{Tr}(A_N A_N^*)^j$ and $x_\lambda := x_1^{r_1} \cdots x_m^{r_m}$. By inserting equation (3.56) into the right-hand side of (3.55) we see that

$$\mu_{2m} = \sum_{\lambda \vdash m} \sum_{\Lambda} b_\Lambda \prod_{\mu \in \Lambda} K_\mu^{\pi_\mu} x_\mu^{\pi_\mu}. \tag{3.61}$$

Similarly, by substituting (3.57) into (3.22) we obtain

$$\mu_{2m} = \sum_{\lambda \vdash m} \sum_{\Lambda} b'_\Lambda \prod_{\mu \in \Lambda} K_\mu^{\pi_\mu} x_\mu^{\pi_\mu}. \tag{3.62}$$

Since the right-hand sides of equations (3.61) and (3.62) identically equal for arbitrary μ, we need to show that $b_\Lambda = b'_\Lambda$

Equations (3.55) and (3.56) give

$$\begin{aligned}
\mu_{2m} &= \frac{(2m-1)!!}{(2\sigma^2)^m} \sum_{\lambda \vdash m} c_\lambda \prod_{j=1}^m \left(\sum_{\mu \vdash j} g_\mu K_\mu(N) x_\mu \right)^{r_j} \\
&= \frac{(2m-1)!!}{(2\sigma^2)^m} \sum_{\lambda \vdash m} c_\lambda \prod_{j=1}^m r_j! \sum_{\substack{\pi_\mu \\ \sum_\mu \pi_\mu = r_j}} \prod_{\mu \vdash j} \frac{g_\mu^{\pi_\mu}}{\pi_\mu !} K_\mu^{\pi_\mu}(N) x_\mu^{\pi_\mu} \\
&= \frac{(2m-1)!!\, m!}{(2\sigma^2)^m} \sum_{\lambda \vdash m} \frac{1}{(1!)^{r_1} \cdots (j!)^{r_j}} \sum_{\Lambda} \prod_{\mu \in \Lambda} \frac{g_\mu^{\pi_\mu}}{\pi_\mu !} K_\mu^{\pi_\mu}(N) x_\mu^{\pi_\mu}.
\end{aligned} \tag{3.63}$$

In the last passage π_μ assumes the same meaning as in equation (3.60), *i.e.* it is the number of repetitions of a partition μ in the union $\lambda = \bigcup_{\mu \in \Lambda} \mu$. Now, let $\lambda = (1^{r_1} \dots m^{r_m})$ and $\mu = (1^{s_1^\mu} \dots j^{s_j^\mu})$ with $\lambda \vdash m$ and $\mu \vdash j$. The frequencies of λ and μ are related by

$$r_k = \sum_{\mu \in \Lambda} s_k^\mu. \tag{3.64}$$

Furthermore, by definition we have

$$g_\mu = \frac{j!}{1^{s_1^\mu} s_1^\mu ! \cdots j^{s_j^\mu} s_j^\mu !}. \tag{3.65}$$

Thus, combining equations (3.63), (3.64) and (3.65) we arrive at

$$\mu_{2m} = \frac{(2m)!}{(2\sigma)^{2m}} \sum_{\lambda \vdash m} \frac{1}{1^{r_1} \cdots m^{r_m}} \sum_{\Lambda} \prod_{j=1}^{m} \prod_{\mu \in \Lambda} \frac{1}{s_j^{\mu}! \pi_\mu!} \prod_{\mu \in \Lambda} K_\mu^{\pi_\mu}(N) x_\mu^{\pi_\mu}. \tag{3.66}$$

Finally, equations (3.21), (3.22), (3.57) and (3.60) give

$$\mu_{2m} = \frac{(2m)!}{(2\sigma)^{2m}} \sum_{\lambda \vdash m} \frac{1}{1^{r_1} \cdots m^{r_m}} \sum_{\Lambda} \prod_{j=1}^{m} \prod_{\mu \in \Lambda} \frac{1}{s_j^{\mu}! \pi_\mu!} \prod_{\mu \in \Lambda} K_\mu'^{\pi_\mu}(N) x_\mu^{\pi_\mu}. \tag{3.67}$$

□

Brouwer and Beenaker [3] computed the leading order asymptotics as $N \to \infty$ of $K_\lambda(N)$. By inserting the right-hand side of (3.57) in equations (3.9) we derive the recursion relations

$$\begin{aligned} NK_{(\lambda_1,\dots,\lambda_k)}(N) &+ \sum_{p+q=\lambda_k} K_{(\lambda_2,\dots,\lambda_k,p,q)}(N) + \sum_{j=1}^{k-1} \lambda_j K_{(\lambda_1,\dots,\lambda_{j-1},\lambda_j+\lambda_k,\lambda_{j+1},\dots,\lambda_{k-1})}(N) \\ &+ \sum_{p+q=\lambda_k} \sum_{j=1}^{k-1} \frac{1}{j!\,(k-j-1)!} \sum_{\sigma \in \mathfrak{S}_{k-1}} K_{(\sigma 1,\dots,\sigma j,p)}(N) K_{(\sigma(j+1),\dots,\sigma(k-1),q)}(N) = 0 \end{aligned} \tag{3.68}$$

with $K_0(N) = 1$. The solution to these equations to leading order is

$$\begin{aligned} K_\lambda(N) &= (-1)^{m+\ell(\lambda)} 2^{\ell(\lambda)} N^{-2m-\ell(\lambda)+2} \\ &\times \frac{(2m+\ell(\lambda)-3)!}{(2m)!} \prod_{j=1}^{m} \frac{\left((2j-1)!\right)^{r_j}}{\left((j-1)!\right)^{2r_j}} + O\left(N^{-2m-\ell(\lambda)}\right). \end{aligned} \tag{3.69}$$

We are now in a position to state the main result of this section.

Theorem 3.9. *We have*

$$\kappa_{2m} = O\left((2m)! N^{-(2-b)(m-1)}\right). \tag{3.70}$$

Proof. This bound follows simply by combining equations (3.69), (3.56) and (3.35). □

4 Proof of the Berry-Esséen inequality

In order to prove the Berry-Esséen bound (2.16), we need an estimate of the radius of convergence of the cumulant expansion (3.48).

Lemma 4.1. *There exists a constant* $\delta > 0$ *such that* $\psi_N(\xi) > 0$ *for* $0 \le |\xi| \le \delta N^{\frac{1-b}{2}}$.

Proof. Since $\psi_N(\xi)$ is entire, the radius of convergence of the Taylor series of $\log \psi_N(\xi)$ is given by the location of the nearest zero to the origin of $\psi_N(\xi)$.

By definition

$$|\psi_N(\xi)| \le \psi_N(0) = \int_{-\infty}^{\infty} f_N(x)dx = 1. \tag{4.1}$$

Suppose that $\psi(\xi)$ has real zeros and let $\bar{\xi}$ be the closest to the origin. Since $\psi_N(\xi)$ is even, we can assume that $\bar{\xi}$ is positive. For $|\xi| < \bar{\xi}$, $0 < \psi_N(\xi) \le 1$, therefore the Taylor series of $\log \psi_N(\xi)$ is convergent in $\left(-\bar{\xi}, \bar{\xi}\right)$. Thus, it also converges in a circle centred at the origin and of radius $\bar{\xi}$. In other words, there are not any complex zero of ψ whose distance from the origin is less than $\bar{\xi}$. Therefore, in the rest of this proof we can take ξ to be real and positive.

A general formula (see [12], p. 514) for moment generating functions gives

$$\left|\psi_N\left(\xi\right) - \sum_{j=0}^{k-1} \left(-1\right)^j \frac{\xi^{2j}}{(2j)!}\mu_{2j}\right| \le \frac{\xi^{2k}}{(2k)!}\mu_{2k}. \tag{4.2}$$

Let us consider the two sums

$$v_{2r-1}(\xi) := \sum_{j=0}^{2r-1} (-1)^j \frac{\xi^{2j}}{(2j)!}\mu_{2j}, \tag{4.3a}$$

$$v_{2s}(\xi) := \sum_{j=0}^{2s} (-1)^j \frac{\xi^{2j}}{(2j)!}\mu_{2j}. \tag{4.3b}$$

Since the Taylor expansion of $\psi_N(\xi)$ is an alternating series, equation (4.2) implies

$$v_{2r-1}(\xi) \le \psi_N(\xi) \le v_{2s}(\xi) \tag{4.4}$$

for any pair of integers $r \ge 1$ and $s \ge 0$. By definition $\mu_2 = 1$, thus the lemma is trivially true for $\xi^2 < 2$.

Let us write

$$\exp\left(-\omega\xi^2/2\right) = w_{2r-1}\left(\xi\right) + u_{2r}\left(\xi\right), \tag{4.5}$$

where

$$w_{2r-1}(\xi) := \sum_{j=0}^{2r-1} (-1)^j \frac{\omega^j \xi^{2j}}{(2j)!}\mu_{2j}^{\mathrm{G}}, \tag{4.6a}$$

$$u_{2r}\left(\xi\right) := \sum_{j=2r}^{\infty} \left(-1\right)^j \frac{\omega^j \xi^{2j}}{(2j)!}\mu_{2j}^{\mathrm{G}}. \tag{4.6b}$$

Recall that $\mu_{2j}^{\mathrm{G}} = (2j-1)!!$ denotes the moments of $\mathcal{N}(0,1)$. We choose $\omega > 4e^2$ and independent of r. We now want to show that for $r \le N$ there exists an appropriate ω such that

$$0 < w_{2r-1}(\xi) \le v_{2r-1}(\xi) \tag{4.7}$$

in the interval

$$\sqrt{2} \le \frac{2e\,(r-1)^{1/2}}{\omega^{1/2}} \le \xi < \frac{2er^{1/2}}{\omega^{1/2}}. \tag{4.8}$$

Since $\omega\xi^2 < 4e^2 r$ the summands in the reminder (4.6b) are strictly decreasing. Therefore, we can write

$$\begin{aligned} w_{2r-1}(\xi) &= \exp\left(-\omega\xi^2/2\right) - \sum_{j=2r}^{\infty} (-1)^j \frac{\omega^j \xi^{2j}}{(2j)!} \mu^{\mathrm{G}}_{2j} \\ &> \exp\left(-\omega\xi^2/2\right) - \frac{\omega^{2r}\xi^{4r}}{(4r)!}\mu^{\mathrm{G}}_{4r} > 0. \end{aligned} \tag{4.9}$$

The last passage is a straightforward consequence of Stirling's formula.

Now, both $w_{2r-1}(\xi)$ and $v_{2r-1}(\xi)$ are alternating sums. Therefore, $w_{2r-1}(\xi) \le v_{2r-1}(\xi)$ if

$$\frac{\mu^{\mathrm{G}}_{2(2j)}}{(4j)!}\omega^{2j}\xi^{4j} - \frac{\mu^{\mathrm{G}}_{2(2j+1)}}{(4j+2)!}\omega^{2j+1}\xi^{4j+2} \le \frac{\mu_{2(2j)}}{(4j)!}\xi^{4j} - \frac{\mu_{2(2j+1)}}{(4j+2)!}\xi^{4j+2} \tag{4.10}$$

for $j \le r-1$. This equation can be rearranged as follows

$$\left(\frac{\xi^2}{2}\right)^{2j}\left(\omega^{2j} - \frac{\mu_{2(2j)}}{\mu^{\mathrm{G}}_{2(2j)}}\right) \le \frac{1}{(2j+1)}\left(\frac{\xi^2}{2}\right)^{2j+1}\left(\omega^{2j+1} - \frac{\mu_{2(2j+1)}}{\mu^{\mathrm{G}}_{2(2j+1)}}\right). \tag{4.11}$$

If we choose $\omega > N^b k$, this inequality holds for $r \le N$ because of proposition 3.5 and equation (3.29). Thus, the statement of the lemma follows if we set $\delta = 2e/\left(N^b\omega\right)^{1/2}$. □

We are now in a position to prove theorem 2.3. From theorem 3.9, we know that for $m \le N$

$$\kappa_{2m} = O\left((2m)!N^{-(2-b)(m-1)}\right). \tag{4.12}$$

Furthermore, from formulae (3.21) and (3.49) it is straightforward to compute the first few cumulants. We have

$$\kappa_2 = 1, \tag{4.13a}$$

$$\kappa_4 = -\frac{3\,\mathrm{Tr}\left(A_N A_N^* - 2\sigma I_N\right)^2}{4\sigma^2 N^3\left(1 - 1/N^2\right)}, \tag{4.13b}$$

$$\kappa_6 = \frac{15\,\mathrm{Tr}\left(A_N A_N^* - 2\sigma I_N\right)^3}{2N^5\sigma^3\left(1-1/N^2\right)\left(1-4/N^2\right)}. \tag{4.13c}$$

Since the cumulant expansion converges up to $\xi \le \delta N^{\frac{1}{2}(1-b)}$, there exists a parameter θ such that

$$\log\psi_N(\xi) = -\frac{\xi^2}{2} + \theta\frac{\xi^4}{N^{2-b}}. \tag{4.14}$$

It turns out that $\theta = O(1)$ as $N \to \infty$. Now, recall that the moment generating function of $\mathcal{N}(0,1)$ is $\psi(\xi) = e^{-\xi^2/2}$. Therefore, we can write

$$|\psi_N(\xi) - \psi(\xi)| = e^{-\xi^2/2}\left|e^{\theta\xi^4/N^{(2-b)}} - 1\right| \le \frac{\theta\xi^4}{N^{2-b}} e^{\theta\xi^4/N^{(2-b)}} e^{-\xi^2/2}, \tag{4.15}$$

where we have used the inequality $|e^z - 1| \le |z|\, e^{|z|}$. The exponential $e^{\theta\xi^4/N^{2-b}}$ is bounded in N provided $\theta = O(1)$. Therefore, the right-hand side of (4.15) becomes

$$|\psi_N(\xi) - \psi(\xi)| \le \frac{C\xi^4}{N^{2-b}} e^{-\xi^2/2}, \tag{4.16}$$

where C can be chosen independent of N.

To complete the proof of equation (4.16), we need to show that if $\xi \le \delta N^{\frac{1}{2}(1-b)}$, then $\theta = O(1)$. Let us write the cumulant expansion as

$$\log\psi_N(\xi) = -\frac{\xi^2}{2} + \frac{\kappa_4\xi^4}{4!} + R_6(N), \tag{4.17}$$

where

$$R_6(N) = \sum_{m=3}^{\infty} \frac{\kappa_{2m}\xi^{2m}}{(2m)!}. \tag{4.18}$$

If a series $\sum_{m=1}^{\infty} c_m$ converges , then $c_m \to 0$ as $m \to \infty$. Therefore, for $m > N$ we must have.

$$\kappa_{2m} = o\Big((2m)!\left(\delta^2 N\right)^{-m(1-b)}\Big), \quad m \to \infty. \tag{4.19}$$

Thus, combining equations (4.12) and (4.19), the reminder (4.18) can be bound by the series

$$\frac{C_1\xi^6}{N^{2(2-b)}} \sum_{m=0}^{N-3} \frac{\xi^{2m}}{N^{(2-b)m}} + \frac{C_2\xi^{2(N+1)}}{N^{(1-b)(N+1)}} \sum_{m=0}^{\infty} \frac{\xi^{2m}}{\left(\delta^* N^{(1-b)}\right)^m}, \tag{4.20}$$

where C_1 and C_2 are constants and $\delta^* > \delta$. For $\xi < N^{\frac{1}{2}(1-b)}$ this sum is $O(1)$ as $N \to \infty$, which implies that θ cannot be an increasing function of N.

Remark 4.2. *There is striking difference between the superexponential rate of convergence discovered by Johansson* [15] *when* A_N *is the identity and the rates of Theorem* 2.1. *Indeed, superexponential rates of convergence to central limit theorems are unusual in probability theory. Theorem* 2.3 *provides some insight into this. When* $A_N = I_N$ *the first* N *moments of* X_N *are gaussian (see equation* (3.47)*) and its first* N *cumulants but* κ_2 *are zero. Therefore, equation* (4.16) *turns into*

$$|\psi_N(\xi) - \psi(\xi)| \le \frac{C\xi^{2(N+1)}}{N^{N+1}} e^{-\xi^2/2}. \tag{4.21}$$

5 Proof of theorem 2.1

5.1 Preliminaries

Let us set $S_N = \delta N^{(1-b)/2}$ and $T_N = N^\gamma$, where $\gamma > 2$. Theorem 2.3 allows us to split the right-hand side of (2.14) as follows:

$$\begin{aligned} e(N) \leq \frac{2C}{N^{2-b}\pi} \int_0^{S_N} \xi^3 e^{-\xi^2/2} d\xi + \frac{2}{\pi S_N} \int_{S_N}^{N^\gamma} e^{-\xi^2/2} d\xi + \frac{2}{\pi S_N} \int_{S_N}^{N^\gamma} \left|\psi_N(\xi)\right| d\xi \\ + \frac{24}{\sqrt{2\pi^3} N^\gamma}. \end{aligned} \tag{5.1}$$

The upper limits of integration can be replaced by infinity. The first integral gives the desired bound. We need to show the remaining terms are of lower order.

The second integral in equation (5.1) can be rewritten as

$$\int_{\delta N^{(1-b)/2}}^{N^\gamma} e^{-\xi^2/2} d\xi \leq \frac{1}{2}\sqrt{\frac{\pi}{2}} \operatorname{erfc}\left(\delta N^{(1-b)/2}/\sqrt{2}\right), \tag{5.2}$$

where

$$\operatorname{erfc}(t) := \frac{2}{\sqrt{2}} \int_t^\infty e^{-x^2} dx \tag{5.3}$$

is the complementary error function. Since $\operatorname{erfc}(t)$ satisfies the inequalities (see, *e.g.* [1], p. 298)

$$\frac{1}{t + \sqrt{t^2+2}} < e^{t^2} \int_t^\infty e^{-\xi^2} d\xi \leq \frac{1}{t + \sqrt{t^2 + \frac{4}{\pi}}}, \tag{5.4}$$

the second integral in (5.1) can be neglected.

The last task that we are left with is to estimate the integral

$$\int_{S_N}^{N^\gamma} \left|\psi_N(\xi)\right| d\xi = O\left(N^{-2}\right). \tag{5.5}$$

5.2 Regularity properties of the distribution of X_N

In general we do not have an explicit formula for $\psi_N(\xi)$ in the interval $(\delta N^{(1-b)/2}, N^\gamma)$. Thus, in order to estimate its behaviour in this range we need to adopt an indirect approach. The idea is to approximate X_N with a random variable X_N^* whose characteristic function allows us to control the third integral in equation (5.1). Then, we will estimate the difference between $e(N)$ and

$$e^*(N) := \sup_{x\in\mathbb{R}} \left|F_N^*(x) - \Phi^*(x)\right|, \tag{5.6}$$

where F_N^* is the approximate distribution function of X_N^* and Φ^* is the distribution function of a random variable close to $\mathcal{N}(0,1)$ (in a sense that will be made precise later).

We first need to discuss some regularity properties of the probability distribution of X_N.[3]

Lemma 5.1. *If* $N > 1$ *the distribution function* F_N *is absolutely continuous, it admits the integral representation*

$$F_N(x) = \int_{-\infty}^{x} f_N(t)dt, \tag{5.7}$$

where $f_N \in L^1(\mathbb{R})$*, is bounded and uniformly continuous. Furthermore,* $\psi_N(\xi) = o\left(\xi^{-1}\right)$ *as* $\xi \to \infty$*.*

Proof. Denote by $\Theta(N)$ the maximal torus of $\mathrm{U}(N)$, *i.e.* the group of diagonal unitary matrices

$$\operatorname{diag}\left(e^{i\theta_1}, \dots, e^{i\theta_N}\right) = \begin{pmatrix} e^{i\theta_1} & & \\ & \ddots & \\ & & e^{i\theta_N} \end{pmatrix}, \tag{5.8}$$

and write $\mathrm{W}(N) = \mathrm{U}(N)/\Theta(N)$. An explicit expression for Haar measure on $\mathrm{U}(N)$ is

$$d\mu_{\mathrm{H}}(U) = \frac{2^{N(N-1)/2}}{(2\pi)^N N!} \prod_{1\le j<k\le N} \sin^2\left(\frac{\theta_k - \theta_j}{2}\right) d\theta_1 \dots d\theta_N d\mu_{\mathrm{W}}, \tag{5.9}$$

where $d\mu_{\mathrm{W}}$ is a normalized Borel measure on $\mathrm{W}(N)$.

Now, recall that

$$X_N = a_1 \cos\theta_1 + \dots + a_N \cos\theta_N, \tag{5.10}$$

where $a_1, \dots, a_N$ are the diagonal elements of A_N. Thus, we can integrate out $d\mu_{\mathrm{W}}$ and study the measure

$$d\mu_{\Theta} = \frac{2^{N(N-1)/2}}{(2\pi)^N N!} \prod_{1\le j<k\le N} \sin^2\left(\frac{\theta_k - \theta_j}{2}\right) d\theta_1 \cdots d\theta_N. \tag{5.11}$$

Since X_N is an absolutely continuous function of $\theta_1, \dots, \theta_N$, if $D \subset [0, 2\pi)^N$ is a set whose image $X_N(D)$ has Lebesgue measure zero, then D must have zero measure too. It follows from equation (5.11) that $\mathbb{P}\left\{X_N \in B\right\} = 0$ for any set B of Lebesgue measure zero. Therefore the probability distribution of X_N is absolutely continuous. Since the only absolutely continuous measures on $\mathbb{R}$ are only those that have a density, F_N admits the integral representation (5.7).

We can say more about f_N. The measure $e^{i\xi X_N} d\mu_\Theta$ is a differential form on the N-dimensional torus. Let

$$\alpha_N = \frac{1}{\sigma} \max_{U\in\mathrm{U}(N)} \left|\operatorname{Tr} A_N U\right| = a_1 + \dots + a_N. \tag{5.12}$$

---

[3] In section 5.3 the ability of estimating a bound for $|f_N(x) - \phi(x)|$ will be essential. Even though $\mathrm{U}(N)$ is compact and Haar measure is absolutely continuous with respect to the Lebesgue measure on $\mathbb{R}^{N^2}$, it is far from obvious that f_N is bounded or even continuous in all $\mathbb{R}$. For example, lemma 5.1 is false for $N = 1$. Indeed, a direct calculation gives $f_1(x) = \left(1 - (x/a)^2\right)^{-1/2}$.

For any neighborhood of $X_N \in [-\alpha_N, \alpha_N]$ we can find a local change of variables that allows us to write

$$e^{i\xi X_N} d\mu_\Theta = e^{i\xi X_N} \mathrm{d}X_N \wedge \omega, \tag{5.13}$$

where ω is $(N-1)$-form on $\Theta(N)$, the symbol $\wedge$ denotes the exterior product and the roman 'd' indicates exterior differentiation.[4] For example, we can choose

$$\begin{aligned} x_N &= a_1 \cos\theta_1 + a_2 \cos\theta_2 + \cdots + a_N \cos\theta_N \\ \varphi_2 &= a_1 \sin(\theta_1 + \beta) + a_2 \sin(\theta_2 - \beta) \\ \varphi_3 &= \theta_3 \\ \vdots\ \ & \quad \vdots \\ \varphi_N &= \theta_N, \end{aligned} \tag{5.14}$$

where β is a real parameter. The Jacobian of this transformation is

$$\begin{aligned} J(x_N, \phi_j) = \frac{\partial(x_N, \varphi_j)}{\partial(\theta_1, \ldots, \theta_N)} &= a_1 a_2 \cos\beta \sin\bigl(\theta_2(x_N, \varphi_j) - \theta_1(x_N, \varphi_j)\bigr) \\ &+ \frac{a_1 a_2 \sin\beta}{2} \cos\bigl(\theta_1(x_N, \phi_j) + \theta_2(x_N, \phi_j)\bigr) \\ &+ \frac{a_1 a_2 \sin\beta}{2} \cos\bigl(\theta_1(x_N, \phi_j) - \theta_2(x_N, \phi_j)\bigr). \end{aligned} \tag{5.15}$$

Thus, the map (5.14) is invertible everywhere except, perhaps, on a surface $\theta_1 = f(\theta_2; \beta)$ where the Jacobian is zero. Appropriate choices of the parameter β in different regions of $\Theta(N)$ allow to define the differential form ω everywhere in $\Theta(N)$. More explicitly, we have

$$\omega = \frac{2^{N(N-1)/2}}{(2\pi)^N N! J(x, \phi_j)} \prod_{1 \le j < k \le N} \sin^2\left(\frac{\theta_k(x, \phi_j) - \theta_j(x, \phi_j)}{2}\right) \mathrm{d}\phi_2 \wedge \cdots \wedge \mathrm{d}\phi_N. \tag{5.16}$$

Let ψ and χ be two differential forms of degrees p and q respectively. The exterior derivative of $\psi \wedge \chi$ is a $(p+q+1)$-form given by

$$\mathrm{d}(\psi \wedge \chi) = \mathrm{d}\psi \wedge \chi + (-1)^p \psi \wedge \mathrm{d}\chi. \tag{5.17}$$

Now, $e^{i\xi X_N}/(i\xi)$ is a 0-form on $\Theta(N)$. Away from the region where $J(x, \phi_j) = 0$ the inverse of the map (5.14) is differentiable with continuous derivatives. Therefore, we have

$$e^{i\xi X_N} d\mu_\Theta = e^{i\xi X_N} \mathrm{d}X_N \wedge \omega = \mathrm{d}\left(\frac{e^{i\xi X_N}}{i\xi}\omega\right) - \frac{e^{i\xi X_N}}{i\xi}\mathrm{d}\omega. \tag{5.18}$$

One can easily verify by direct calculation that ω is not closed, *i.e.* $\mathrm{d}\omega \neq 0$.

---

[4] While an absolute continuous measure $d\mu$ on a smooth manifold can always be interpreted as a differential form, it does not mean that it is the exterior derivative of another form. Indeed, $d\mu_\Theta$ is not. We use the notation 'd' to emphasise this difference, because it is important in what follows.

If ψ is a differential form of degree p and Ω is a manifold of dimension $p+1$, then Stokes' theorem states that

$$\int_\Omega \mathrm{d}\psi = \int_{\partial\Omega} \psi, \tag{5.19}$$

where $\partial\Omega$ denotes the boundary of Ω. An N-dimensional torus is a compact manifold without boundary, therefore the right-hand side of (5.19) is zero. As a consequence, integrating both sides of equation (5.18) we obtain

$$\psi_N(\xi) = \int_{\Theta(N)} e^{i\xi X_N} \left(\mathrm{d}X_N \wedge \omega\right) = -\frac{1}{i\xi}\int_{\Theta(N)} e^{i\xi X_N} \mathrm{d}\omega. \tag{5.20}$$

It follows from the Riemann-Lebesgue lemma that $\psi_N(\xi) = o\left(\xi^{-1}\right)$ as $\xi \to \infty$ and is integrable. Thus, the inverse Fourier transform

$$f_N(x) = \frac{1}{2\pi}\int_{-\infty}^{\infty} e^{-i\xi x}\psi_N(\xi)d\xi \tag{5.21}$$

is well defined, bounded and uniformly continuous. □

5.3 Smoothing

From the discussion in §5.1, it follows that a necessary (not sufficient) condition for $\int_{S_N}^{N^\gamma} \left|\psi_N\left(\xi\right)\right| d\xi$ to decay fast enough is $\psi_N(\xi) = O\left(\xi^{-4}\right)$ as $\xi \to \infty$. Indeed, if A_N is of full rank and its spectrum is not degenerate, equation (2.8) and the asymptotic formula

$$J_k(x) \sim \sqrt{\frac{2}{\pi x}}\cos\left(x - \tfrac{1}{4}k\pi - \tfrac{1}{4}\pi\right), \quad x \to \infty \tag{5.22}$$

imply that $\psi_N\left(\xi\right) = O\bigl(\xi^{-N^2/2}\bigr)$. Therefore, f_N has continuous derivatives at least up to order $N^2/2 - 2$. In other words, f_N becomes increasingly smooth as N grows.

If a function has continuous derivatives of order p, then its Fourier transform is $o\left(\xi^{-p}\right)$ as $\xi \to \infty$. This suggests smoothing F_N with an appropriate test function. More precisely, we define

$$F_N^*(x) := \left[F_N * \chi_\epsilon\right](x) = \int_{-\infty}^{\infty} F_N(t)\chi_\epsilon(x-t)dt, \tag{5.23}$$

where $\chi_\epsilon \in \mathcal{C}_0^\infty\left(\mathbb{R}\right)$ and is normalized to one. Our choice will be the test function

$$\chi_\epsilon(x) := \begin{cases} \frac{1}{g\epsilon}\exp\left(-\frac{1}{1-\left(\frac{x}{\epsilon}\right)^2}\right) & \text{if } x \in (-\epsilon,\epsilon), \\ 0 & \text{if } x \in \mathbb{R}\setminus(-\epsilon,\epsilon), \end{cases} \tag{5.24}$$

where

$$g := \int_{-1}^{1} \exp\left(-\frac{1}{1-x^2}\right) dx = 0.44399\ldots \tag{5.25}$$

By differentiating F_N^* and integrating by parts we obtain

$$f_N^*(x) := F_N^{*\prime}(x) = \left[f_N * \chi_\epsilon\right](x) = \int_{-\infty}^{\infty} f_N(t)\chi_\epsilon(x-t)dt. \tag{5.26}$$

The convolution f_N^* is positive and

$$\int_{-\infty}^{\infty} f_N^*(x)\,dx = \left(\int_{-\infty}^{\infty} f_N(x)dx\right)\left(\int_{-\infty}^{\infty} \chi_\epsilon(x)\,dx\right) = 1. \tag{5.27}$$

In addition, $f_N^* \in \mathcal{C}_0^\infty(\mathbb{R})$ too and

$$\psi_N^*(\xi) := \int_{-\infty}^{\infty} e^{i\xi x} f_N^*(x)dx = \psi_N(\xi)\hat{\chi}_\epsilon(\xi), \tag{5.28}$$

where

$$\hat{\chi}_\epsilon(\xi) := \int_{-\infty}^{\infty} e^{i\xi x}\chi_\epsilon(x)\,dx. \tag{5.29}$$

Let us introduce

$$\Delta(x) := F_N(x) - \Phi(x), \qquad \Delta^*(x) := \left[\Delta * \chi_\epsilon\right](x). \tag{5.30}$$

Then, we write

$$e(N) = \max_{x\in\mathbb{R}} \left|\Delta(x)\right|, \qquad e^*(N) = \max_{x\in\mathbb{R}} \left|\Delta^*(x)\right|. \tag{5.31}$$

Formula (2.14) still holds if we replace $\psi_N(\xi)$ and $\psi(\xi)$ with $\psi_N^*(\xi)$ and $\psi^*(\xi) := \psi(\xi)\hat{\chi}_\epsilon(\xi)$ respectively. Indeed, let $S_N = \delta N^{(1-b)/2}$ and $T_N = N^\gamma$ with $\gamma > 2$. We have

$$\begin{aligned} e^*(N) \le \frac{2}{\pi}\int_0^{S_N} \left|\frac{\psi_N^*(\xi)-\psi^*(\xi)}{\xi}\right| d\xi + \frac{2}{\pi S_N}\int_{S_N}^{N^\gamma} \left|\psi_N^*(\xi)\right| d\xi \\ + \frac{2}{\pi S_N}\int_{S_N}^{N^\gamma} \left|\psi^*(\xi)\right| d\xi + \frac{24m}{\pi N^\gamma}, \end{aligned} \tag{5.32}$$

where

$$m := \max_{\mathbb{R}} \left|\phi * \chi_\epsilon\right|. \tag{5.33}$$

Now,

$$\left|\psi_N^* - \psi^*\right| = \left|\psi_N - \psi\right|\left|\hat{\chi}_\epsilon\right| \le \left|\psi_N - \psi\right|, \tag{5.34}$$

where we have used $|\hat{\chi}_\epsilon| \le 1$, which holds for any characteristic function. Therefore the Berry-Esséen inequality (2.16) applies to $|\psi_N^* - \psi^*|$ too and

$$\frac{2}{\pi}\int_0^{S_N} \left|\frac{\psi_N^*(\xi)-\psi^*(\xi)}{\xi}\right| d\xi = O\left(N^{2-b}\right). \tag{5.35}$$

Equation (5.2) gives

$$\int_{S_N}^{N^\gamma} |\psi^*(\xi)|\, d\xi \le \int_{S_N}^{N^\gamma} e^{-\xi^2/2} d\xi \le \frac{1}{2}\sqrt{\frac{\pi}{2}}\operatorname{erfc}\left(\delta N^{(1-b)/2}/\sqrt{2}\right). \tag{5.36}$$

In order to complete the proof of equation (2.12), we need to show that, for appropriate choices of the smoothing parameter ϵ, $e(N) \le Ce^*(N)$, and that the integral

$$\int_{S_N}^{N^\gamma} |\psi_N^*(\xi)|\, d\xi = \int_{S_N}^{N^\gamma} |\psi_N(\xi)|\, |\hat{\chi}_\epsilon(\xi)|\, d\xi \tag{5.37}$$

is sufficiently small. The appropriate choice of ϵ for which these two statements are true is a delicate balance. As ϵ decreases $e^*(N)$ will approach $e(N)$. However, if the support of χ_ϵ is too small, its Fourier transform might spread for a range of $\xi > S_N$ large enough to prevent the integral (5.37) from decaying at a sufficiently fast rate.

The leading order asymptotics of $\hat{\chi}_\epsilon(\xi)$ can be computed using the method of steepest descent. We report the calculation in the appendix. We have

$$\begin{aligned}\hat{\chi}_\epsilon(\xi) &= \frac{2}{g(\epsilon\xi)^{3/4}}\sqrt{\frac{\pi}{\sqrt{2}}}\cos\left(\epsilon\xi - (\epsilon\xi)^{1/2} - \tfrac{3}{8}\pi\right)\\ &\quad\times \exp\left(-(\epsilon\xi)^{1/2} - \tfrac{1}{4}\right)\left(1 + O\left((\epsilon\xi)^{-1/2}\right)\right), \quad \epsilon\xi \to \infty.\end{aligned} \tag{5.38}$$

For this approximation to be meaningful $\xi > 1/\epsilon$. Therefore, we cannot choose $\epsilon < C/S_N$, otherwise the bound on the decay rate of the integral (5.37) would not be adequate.

It remains to establish if $\epsilon = O\left(S_N^{-1}\right)$ leads to a good enough approximation to $e(N)$. In order not to loose information on the behaviour of $\Delta(x)$, the smoothing parameter needs to be comparable with the rate of oscillation of $\Delta(x)$. In other words, we need a bound on $|\Delta'(x)|$. Such a bound can be obtained, once again, using the Berry-Esséen inequality (2.16):

$$\begin{aligned}|\Delta'(x)| = |f_N(x) - \phi(x)| &\le \int_{-S_N}^{S_N} |\psi_N(\xi) - \psi(\xi)|\, d\xi\\ &\quad + 2\int_{S_N}^{\infty} |\psi_N(\xi)|\, d\xi + 2\int_{S_N}^{\infty} |\psi(\xi)|\, d\xi.\end{aligned} \tag{5.39}$$

By comparing this inequality with (5.1), we see that the first and third integral are $O\left(N^{-(2-b)}\right)$; the second integral might possibly be bigger by a factor πS_N. It follows that

$$|\Delta'(x)| \le CS_N e(N). \tag{5.40}$$

This is sufficient for our purposes. The following lemma completes the proof of theorem 2.1.

Lemma 5.2. *Suppose that* $|\Delta'(x)| \le e(N)/\eta_N$. *Then, for* $\epsilon = \eta_N$ *there exists a positive constant* $C = O(1)$ *such that* $e(N) \le Ce^*(N)$.

Proof. Since $\Delta(t)$ is continuous and $-1 \leq \Delta(t) \leq 1$, there exists a $t_0 \in \mathbb{R}$ such that $e(N) = \Delta(t_0)$. Then, we have

$$\Delta(t_0 + y) \geq e(N)\left(1 - \frac{y}{\eta_N}\right), \quad \text{for } y > 0. \tag{5.41}$$

Now set

$$x = t_0 + r\eta_N, \quad t = r\eta_N - y, \tag{5.42}$$

where $0 < r < 1$ is a parameter whose exact value is to be determined. Equation (5.41) becomes

$$\Delta\left(x - t\right) \geq e(N)\left(1 - r + \frac{t}{\eta_N}\right), \quad \text{for } |t| \leq r\eta_N. \tag{5.43}$$

Substituting this bound into the definition

$$\Delta^*(x) = \int_{-\infty}^{\infty} \Delta(x-t)\chi_{\eta_N}(t)dt \tag{5.44}$$

we arrive at

$$e^*\left(N\right) \geq \Delta^*(x) \geq e(N)\left(1 - r\right)\int_{|t|\leq r\eta_N} \chi_{\eta_N}(t)dt - e(N)\int_{|t|\geq r\eta_N} \chi_{\eta_N}(t)dt, \tag{5.45}$$

where we have used the inequality $\Delta(t) \geq -e(N) \geq -1$ and the fact that the linear term in (5.43) does not contribute because $\chi_{\eta_N}(t)$ is even.

Now, it turns out that the two integrals in (5.45) are independent of η_N. Indeed,

$$\int_{|t|\geq r\eta_N} \chi_{\eta_N}(t)dt = \frac{2}{g}\int_r^1 \exp\left(-\frac{1}{1-t^2}\right)dt. \tag{5.46}$$

Define

$$h(r) := r + \frac{2}{g}\left(2 - r\right)\int_r^1 \exp\left(-\frac{1}{1-t^2}\right)dt. \tag{5.47}$$

Equation (5.45) can now be rewritten

$$e^*(N) \geq e(N)\bigl(1 - h(r)\bigr). \tag{5.48}$$

Since $h(r)$ has a minimum near $2/3$ and $h(2/3) = 0.77646.. \approx 4/5$, the statement of the lemma follows with $C = 5$. ☐

We achieve the rate of convergence in equation (2.12) if we set

$$\eta_N = \frac{1}{CS_N} = \frac{1}{C\delta N^{(1-b)/2}}. \tag{5.49}$$

6 Proof of theorem 2.2

We need to prove equation (2.13).

Let us choose the parameter η_N in lemma 5.2 to be

$$\eta_N = N^{-\zeta}, \tag{6.1}$$

where $0 < \zeta < \frac{1}{2}(1-b)$. Equation (5.39) gives

$$|f_N(x) - \phi(x)| \leq C_\zeta N^{-2+b+\zeta} \tag{6.2}$$

for some constant C_ζ.

Now take $\zeta' > 0$ and write

$$\begin{aligned}\int_{-\infty}^{\infty} |f_N(x) - \phi(x)|\, dx &\leq 2\int_0^{N^{\zeta'}} |f_N(x) - \phi(x)|\, dx \\ &\quad + 2\int_{N^{\zeta'}}^{\infty} f_N(x)dx + 2\int_{N^{\zeta'}}^{\infty} \phi(x)dx \\ &\leq C_{\zeta,\zeta'} N^{-2+b+\zeta+\zeta'} + 2\int_{N^{\zeta'}}^{\infty} f_N(x)dx + 2\int_{N^{\zeta'}}^{\infty} \phi(x)dx.\end{aligned} \tag{6.3}$$

We have

$$\begin{aligned}2\int_{N^{\zeta'}}^{\infty} f_N(x)dx &= \int_{-\infty}^{-N^{\zeta'}} f_N(x)dx + \int_{N^{\zeta'}}^{\infty} f_N(x)dx \\ &= F_N\left(-N^{\zeta'}\right) + 1 - F_N\left(N^{\zeta'}\right).\end{aligned} \tag{6.4}$$

Similarly

$$\begin{aligned}2\int_{N^{\zeta'}}^{\infty} \phi(x)dx &= \int_{-\infty}^{-N^{\zeta'}} \phi(x)dx + \int_{N^{\zeta'}}^{\infty} \phi(x)dx \\ &= \Phi\left(-N^{\zeta'}\right) + 1 - \Phi\left(N^{\zeta'}\right).\end{aligned} \tag{6.5}$$

Therefore, rearranging the terms and using the identity $1 - \Phi(x) = \Phi(-x)$ we obtain

$$\begin{aligned}2\int_{N^{\zeta'}}^{\infty} f_N(x)dx + 2\int_{N^{\zeta'}}^{\infty} \phi(x)dx &= \Big(F_N\left(-N^{\zeta'}\right) - \Phi\left(-N^{\zeta'}\right)\Big) \\ &\quad - \Big(F_N\left(N^{\zeta'}\right) - \Phi\left(N^{\zeta'}\right)\Big) + 4\Big(1 - \Phi\left(N^{\zeta'}\right)\Big).\end{aligned} \tag{6.6}$$

Finally, equation (2.13) follows from (2.12) and (5.4) by setting $\epsilon = \zeta + \zeta'$.

Acknowledgements

We would like to express our gratitude to Paul Bourgade, Oliver Johnson and Roman Schubert and for helpful discussions. While this research was carried out, F. Mezzadri was partially supported by EPSRC grant no. EP/G019843/1 and by a Leverhulme Research Fellowship.

Appendix. The leading order asymptotics of $\hat{\chi}_\epsilon(\xi)$

The purpose of this appendix is to compute an explicit formula for the leading order asymptotics of

$$\hat{\chi}_\epsilon\left(\xi\right) := \int_{-\infty}^{\infty} e^{i\xi x}\chi_\epsilon(x)dx = \frac{1}{g\epsilon}\int_{-\epsilon}^{\epsilon} \exp\left(i\xi x - \frac{1}{1-\left(\frac{x}{\epsilon}\right)^2}\right)dx \tag{A.7}$$

in the limit $\xi \to \infty$. Since $\hat{\chi}_\epsilon(\xi) = \hat{\chi}_1\left(\epsilon\xi\right)$, for the sake of simplicity we set $\epsilon = 1$ and write $\hat{\chi} = \hat{\chi}_1$. More explicitly, we study the integral

$$\hat{\chi}\left(\xi\right) = \frac{2}{g}\operatorname{Re}\int_0^1 \exp\left(i\xi x - \frac{1}{1-x^2}\right)dx. \tag{A.8}$$

Proposition. *We have*

$$\begin{aligned}\hat{\chi}\left(\xi\right) = {} & \frac{2}{g\xi^{3/4}}\sqrt{\frac{\pi}{\sqrt{2}}}\cos\left(\xi - \xi^{1/2} - \tfrac{3}{8}\pi\right) \\ & \times \exp\left(-\xi^{1/2} - \tfrac{1}{4}\right)\left(1 + O\left(\xi^{-1/2}\right)\right), \quad \xi \to \infty.\end{aligned} \tag{A.9}$$

Proof. (A.8) can be estimated using the method of steepest descents. The integrand is not analytic at one, so we look at

$$\frac{2}{g}\operatorname{Re}\int_0^{1-\delta} \exp\left(i\xi x - \frac{1}{1-x^2}\right)dx, \tag{A.10}$$

where $\delta > 0$ is small. The difference between (A.8) and (A.10) is bounded by $2\delta e^{-1/(2\delta)}/g$.

Since we are interested only in the real part of (A.10), for large ξ the origin will not contribute to leading order; the main contribution should come from a small neighbourhood near one.

Consider the argument of the exponential:

$$f(x) := i\xi x - \frac{1}{1-x^2}. \tag{A.11}$$

Its saddle points are the solutions of the equation

$$i\xi\left(1-x^2\right)^2 - 2x = 0. \tag{A.12}$$

For large ξ the roots of this polynomial can be computed perturbatively in the parameter $1/\xi$. In other words, we look for a solution near one with an asymptotic expansion of the form

$$\bar{x}(\xi) = 1 + \frac{x_1}{\xi^{\alpha}} + \frac{x_2}{\xi^{2\alpha}} + O\left(\xi^{-3\alpha}\right), \quad \xi \to \infty, \tag{A.13}$$

where α is a rational power. By substituting this expression into (A.12), one finds that the two sides of the equation can be balanced only if $\alpha = 1/2$ and that the first two coefficients are

$$x_1 = \pm\frac{e^{\frac{3}{4}\pi i}}{\sqrt{2}} \quad \text{and} \quad x_2 = 0. \tag{A.14}$$


Im(x)
C
$\times\ 1 + \frac{e^{\frac{3}{4}\pi i}}{\sqrt{2}}\frac{1}{\xi^{1/2}}$
0
$1-\delta$
Re(x)
$\times\ 1 - \frac{e^{\frac{3}{4}\pi i}}{\sqrt{2}}\frac{1}{\xi^{1/2}}$

Figure 1: The deformation of the interval $[0, 1-\delta]$ and the saddle points of $f(x)$.

Provided δ is sufficiently small, the interval of integration of (A.10) can be deformed into a contour asymptotically equivalent to the steepest descent path passing through

$$\bar{x}(\xi) = 1 + \frac{e^{\frac{3}{4}\pi i}}{\sqrt{2}}\frac{1}{\xi^{1/2}} + O\left(\xi^{-3/2}\right), \quad \xi \to \infty. \tag{A.15}$$

(See figure 1.) Such a deformation is not possible for the critical point

$$1 - \frac{e^{\frac{3}{4}\pi i}}{\sqrt{2}}\frac{1}{\xi^{1/2}} + O\left(\xi^{-3/2}\right), \quad \xi \to \infty. \tag{A.16}$$

Trivial algebra gives

$$f\big(\bar{x}(\xi)\big) = i\left(\xi - \xi^{1/2}\right) - \xi^{1/2} - \frac{1}{4} + O\left(\xi^{-1/2}\right), \quad \xi \to \infty, \tag{A.17a}$$

$$f''\big(\bar{x}(\xi)\big) = 2\sqrt{2}e^{-i\frac{\pi}{4}}\xi^{3/2} + O\left(\xi^{1/2}\right), \quad \xi \to \infty. \tag{A.17b}$$

The tangent to the steepest descent path at $\bar{x}$ has equation

$$x(t) = \bar{x}(\xi) + te^{-\frac{3}{8}\pi i}. \tag{A.18}$$

Therefore, we have

$$\begin{aligned}\int_0^{1-\delta} \exp\left(i\xi x - \frac{1}{1-x^2}\right) dx &= \int_C \exp\left(i\xi x - \frac{1}{1-x^2}\right) dx \\ &\sim e^{-\frac{3}{8}\pi i}\sqrt{\frac{2\pi}{\left|f''\left(\bar{x}(\xi)\right)\right|}}\exp\Big(f\left(\bar{x}(\xi)\right)\Big), \quad \xi \to \infty.\end{aligned} \tag{A.19}$$

Finally, by inserting equations (A.17) into (A.19) we arrive at (A.9), provided $\delta < \xi^{-\beta}$ and $\beta > 3/4$.

□

References

[1] Abramowitz M. and Stegun I. *Handbook of Mathematical Functions.* United States Department of Commerce, National Bureau of Standards, Applied Mathematics Series 55, Washington, 1972.

[2] Bars I. U(N) integral for the generating function in lattice gauge theory. *J. Math. Phys.* **21** (1980), 2678–2681.

[3] Brouwer P. W. and Beenakker W. J. Diagrammatic method of integration over the unitary group, with applications to quantum transport mesoscopic systems. *J. Math. Phys.* **37** (1996), 4904–4934.

[4] Brower R. C., Rossi P. and Tan C.-I. The external field problem for QCD. *Nucl. Phys. B* **190** (1981), 699–718.

[5] Chatterjee S. and Meckes E. Multivariate normal approximation using exchangeable pairs. *ALEA Lat. Am. J. Prob. Math. Stat.* **4** (2008), 257–283.

[6] Collins B. and Stolz M. Borel theorems for random matrices from the classical compact symmetric spaces. *Ann. Prob.* **36** (2008), 876–895.

[7] Damgaard P. H. and Splittorff. Spectral sum rules of the Dirac operator and partially quenched chiral condensates. *Nucl. Phys. B* **572** (2000), 478–498.

[8] De Wit B. and 't Hooft G. Nonconvergence of the $1/N$ expansion for SU(N) gauge fields on a lattice. *Phys. Lett. B* **69** (1977), 61–64.

[9] Diaconis P. and Shahshahani M. On the eigenvalues of random matrices. Studies in applied probability. *J. Appl. Prob.* **31A** (1994), 49–62.

[10] D'Aristotile A., Diaconis P. and Newman C. Brownian motion and the classical groups. *Probability, Statistics and their Applications: Papers in Honor of Rabi Bhattacharya.* IMS Lect. Notes Monogr. Ser. **41** (2003), 97–116.

[11] Diaconis P. and Evans S. N. Linear functionals of eigenvalues of random matrices. *Trans. Amer. Math. Soc.* **353** (2001), 2615–2633.

[12] Feller W. *An Introduction to Probability Theory and Its Applications.* Vol. II, second edition. John Wiley & Sons, Inc., New York, 1970.

[13] Haake F., Kuś M., Sommers H.-J., Schomerus H. and Życzkowski K. Secular determinants of random unitary matrices. *J. Phys. A: Math. Gen.* **29** (1996), 3641–3658.

[14] Hughes C. P. and Rudnick Z. Mock-Gaussian behaviour for linear statistics of classical compact groups. Random matrix theory. *J. Phys. A: Math. Gen.* **36** (2003), 2919–2932.

[15] Johansson K. On random matrices from the classical compact group. *Ann. Math. (2)* **145** (1997), 519–545.

[16] Macdonald I. G. *Symmetric functions and Hall polynomials.* Second edition. Oxford University Press, New York, 1995.

[17] Meckes E. *An Infinitesimal Version of Stein's Method of Exchangeable Pairs.* PhD Thesis, Stanford University, 2006.

[18] Meckes E. Linear functions on the classical matrix groups. *Trans. Amer. Math. Soc.* **360** (2008), 5355–5366.

[19] Moiseyev N. Quantum theory of resonances: calculating energies, widths and cross-sections by complex scaling. *Phys. Rep.* **302** (1998), 212–293.

[20] Okołowicz J., Płoszajczak M. and Rotter I. Dynamics of quantum systems embedded in a continuum. *Phys. Rep.* **374** (2003), 271–383.

[21] Pastur L. and Vasilchuk V. On the moments of traces of classical groups. *Commun. Math. Phys.* **252** (2004), 149–166.

[22] Samuel S. U(N) integrals, $1/N$, and the De Wit-'t Hooft anomalies. *J. Math. Phys.* **21** (1980), 2695–2703.

[23] Schlittgen B. and Wettig T. Generalizations of some integrals over the unitary group. *J. Phys. A: Math. Gen.* **36** (2003), 3195–3201.

[24] Soshnikov A. The central limit theorem for local linear statistics in the classical compact groups and related combinatorial identities. *Ann. Prob.* **28** (2000), 1353–1370.

[25] Stolz M. On the Diaconis-Shahshahani method in random matrix theory. *J. Algeb. Comb.* **22** (2005), 471–491.

Department of Mathematics
University of Bristol
Bristol BS8 1TW, UK
Email: `j.p.keating@bristol.ac.uk`
Email: `f.mezzadri@bristol.ac.uk`
Email: `b.singphu@bristol.ac.uk`

16 November 2010